

Predictors of outcome in patients with severe sepsis or septic shock due to extended-spectrum  $\beta$ -lactamase-producing Enterobacteriaceae



A. Russo , M. Falcone , B. Gutiérrez-Gutiérrez , E. Calbo ,  
B. Almirante , P.L. Viale , A. Oliver , P. Ruiz-Garbajosa ,  
O. Gasch , M. Gozalo , J. Pitout , M. Akova , C. Peña ,  
J.M. Cisneros , A. Hernández-Torres , A. Farcomeni , N. Prim ,  
J. Origüen , G. Bou , E. Tacconelli , M. Tumbarello ,  
A. Hamprecht , I. Karaïskos , C. de la Calle , F. Pérez ,  
M.J. Schwaber , J. Bermejo , W. Lowman , P.-R. Hsueh ,  
M. Mora-Rillo , J. Rodríguez-Gomez , M. Souli , R.A. Bonomo ,  
D.L. Paterson , Y. Carmeli , A. Pascual , J. Rodríguez-Baño ,  
M. Venditti , REIPI/ESGBIS/INCREMENT investigators

PII: S0924-8579(18)30186-9  
DOI: [10.1016/j.ijantimicag.2018.06.018](https://doi.org/10.1016/j.ijantimicag.2018.06.018)  
Reference: ANTAGE 5476

To appear in: *International Journal of Antimicrobial Agents*

Received date: 13 April 2018  
Revised date: 6 June 2018  
Accepted date: 18 June 2018

Please cite this article as: A. Russo , M. Falcone , B. Gutiérrez-Gutiérrez , E. Calbo ,  
B. Almirante , P.L. Viale , A. Oliver , P. Ruiz-Garbajosa , O. Gasch , M. Gozalo , J. Pitout ,  
M. Akova , C. Peña , J.M. Cisneros , A. Hernández-Torres , A. Farcomeni , N. Prim , J. Origüen ,  
G. Bou , E. Tacconelli , M. Tumbarello , A. Hamprecht , I. Karaïskos , C. de la Calle , F. Pérez ,  
M.J. Schwaber , J. Bermejo , W. Lowman , P.-R. Hsueh , M. Mora-Rillo , J. Rodríguez-Gomez ,  
M. Souli , R.A. Bonomo , D.L. Paterson , Y. Carmeli , A. Pascual , J. Rodríguez-Baño ,  
M. Venditti , REIPI/ESGBIS/INCREMENT investigators, Predictors of outcome in patients with  
severe sepsis or septic shock due to extended-spectrum  $\beta$ -lactamase-producing Enterobacteriaceae,  
*International Journal of Antimicrobial Agents* (2018), doi: [10.1016/j.ijantimicag.2018.06.018](https://doi.org/10.1016/j.ijantimicag.2018.06.018)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## 1 HIGHLIGHTS

- 2 - Few data are reported in literature regarding sepsis or septic shock due to ESBL strains.
- 3 - The INCREMENT project is the largest international cohort study on ESBL infections.
- 4 - Septic shock due to ESBL strains is associated with high 30-day mortality rate.
- 5 - Escalation of antibiotic therapy is an important determinant of mortality.
- 6 - Carbapenems should be used only in strains with resistance to  $\beta$ -lactam/ $\beta$ -lactamase inhibitors

7

# **Predictors of outcome in patients with severe sepsis or septic shock due to extended-spectrum $\beta$ -lactamase-producing Enterobacteriaceae**

A. Russo<sup>1</sup>, M. Falcone<sup>1</sup>, B. Gutiérrez-Gutiérrez<sup>2</sup>, E. Calbo<sup>3</sup>,  
B. Almirante<sup>4</sup>, P.L. Viale<sup>5</sup>, A. Oliver<sup>6</sup>, P. Ruiz-Garbajosa<sup>7</sup>, O. Gasch<sup>8</sup>, M. Gozalo<sup>9</sup>, J.  
Pitout<sup>10</sup>, M. Akova<sup>11</sup>, C. Peña<sup>12</sup>, J.M. Cisneros<sup>13</sup>, A. Hernández-Torres<sup>14</sup>, A. Farcomeni<sup>1</sup>,  
N. Prim<sup>15</sup>, J. Origüen<sup>16</sup>, G. Bou<sup>17</sup>, E. Tacconelli<sup>18</sup>, M. Tumbarello<sup>19</sup>, A. Hamprecht<sup>20</sup>, I.  
Karaiskos<sup>21</sup>, C. de la Calle<sup>22</sup>, F. Pérez<sup>23</sup>, M.J. Schwaber<sup>24</sup>, J. Bermejo<sup>25</sup>, W. Lowman<sup>26</sup>,  
P.-R. Hsueh<sup>27</sup>, M. Mora-Rillo<sup>28</sup>, J. Rodríguez-Gómez<sup>29</sup>, M. Souli<sup>30</sup>, R.A. Bonomo<sup>23,31</sup>,  
D.L. Paterson<sup>32</sup>, Y. Carmeli<sup>24</sup>, A. Pascual<sup>2</sup>, J. Rodríguez-Baño<sup>2</sup>, M. Venditti<sup>1</sup> and  
REIPI/ESGBIS/INCREMENT investigators.

<sup>1</sup>*Department of Public Health and Infectious Diseases, Policlinico Umberto I, University of Rome La Sapienza, Rome, Italy*

<sup>2</sup>*Unidad Clínica de Enfermedades Infecciosas y Microbiología, Hospital Universitario Virgen Macarena / Universidad de Sevilla / Instituto de Biomedicina de Sevilla, Seville, Spain*

<sup>3</sup>*Hospital Universitari Mútua de Terrassa, Barcelona, Spain*

<sup>4</sup>*Hospital Universitari Vall d'Hebrón, Barcelona, Spain*

<sup>5</sup>*Teaching Hospital Policlinico S. Orsola Malpighi, Bologna, Italy*

<sup>6</sup>*Hospital Universitario Son Espases, Instituto de Investigación Illes Balears (IdISBa), Palma de Mallorca, Spain*

<sup>7</sup>*Hospital Ramón y Cajal, Madrid, Spain.*

<sup>8</sup>*Corporació Sanitària Parc Taulí, Sabadell, Barcelona, Spain*

<sup>9</sup>*Hospital Universitario Marqués de Valdecilla - IDIVAL, Santander, Spain*

<sup>10</sup>*Department of Pathology and Laboratory Medicine, University of Calgary, Calgary, Alberta, Canada*

<sup>11</sup>*Hacettepe University School of Medicine, Ankara, Turkey*

<sup>12</sup>*Hospital Universitari de Bellvitge, Barcelona, Spain*

<sup>13</sup>*Infectious Diseases, Microbiology and Preventive Medicine, Institute of Biomedicine of Seville (IBiS), University Hospital Virgen del Rocío, CSIC, University of Seville, Seville, Spain.*

<sup>14</sup>Hospital Universitario Virgen de la Arrixaca, Murcia, Spain

<sup>15</sup>Hospital de la Santa Creu i Sant Pau, Barcelona, Spain

<sup>16</sup>Hospital Universitario 12 de Octubre, Madrid, Spain.

<sup>17</sup>Complejo Hospitalario Universitario A Coruña, A Coruña, Spain.

<sup>18</sup>Universitätsklinikum Tübingen, Tübingen, Germany

<sup>19</sup>Catholic University of the Sacred Heart, Rome, Italy

<sup>20</sup>Institut für Mikrobiologie, Immunologie und Hygiene Universitätsklinikum Köln,  
Cologne, Germany

<sup>21</sup>Hygeia General Hospital, Athens, Greece

<sup>22</sup>Hospital Clinic, Barcelona, Spain

<sup>23</sup>Research Service, Louis Stokes Cleveland Department of Veterans Affairs Medical  
Center, Cleveland, Ohio, USA

<sup>24</sup> Division of Epidemiology and Preventive Medicine, Tel Aviv Sourasky Medical  
Center, Tel Aviv, Israel, and National Center for Infection Control, Israel Ministry of  
Health, Tel Aviv, Israel, affiliated with Sackler Faculty of Medicine, Tel Aviv  
University, Tel Aviv, Israel

<sup>25</sup>Hospital Español, Rosario, Argentina

<sup>26</sup>Wits Donald Gordon Medical Centre, Johannesburg, South Africa.

<sup>27</sup>National Taiwan University Hospital, National Taiwan University Hospital College of  
Medicine, Taipei, Taiwan.

<sup>28</sup>Hospital Universitario La Paz-IdiPAZ, Madrid, Spain.

<sup>29</sup> Intensive Care Unit, Biomedical Research Institute of Cordoba (IMIBIC)/ Reina Sofia  
University Hospital/University of Cordoba, Córdoba, Spain

<sup>30</sup>University General Hospital Attikon, Chaidari, Greece

<sup>31</sup>Departments of Medicine, Pharmacology, Biochemistry, Molecular Biology and  
Microbiology, Case Western Reserve University School of Medicine, Cleveland Ohio,  
USA

<sup>32</sup>University of Queensland Centre for Clinical Research, The University of  
Queensland, Herston, Brisbane, Australia

**Word count:** 2791

**Running head:** septic shock due to ESBL infections.

**Corresponding author:**

Prof. Mario Venditti

Department of Public Health and Infectious Diseases

Policlinico Umberto I, Viale dell'Università 37, 00161 "Sapienza" University of Rome,

Rome, Italy

email: [mario.venditti@uniroma1.it](mailto:mario.venditti@uniroma1.it)

Tel/fax: 003906491749

99 **ABSTRACT**

100 **Purpose:** Few data are reported in literature regarding severe sepsis or septic shock due  
101 to extended-spectrum  $\beta$ -lactamases (ESBL)-producing Enterobacteriaceae (E). The aim  
102 of this study was to assess predictors of outcome in septic patients with bloodstream  
103 infection (BSI) caused by ESBL-E.

104 **Methods:** Patients with severe sepsis or septic shock and BSI due to ESBL-E were  
105 selected from the INCREMENT database. The primary endpoint of the study was the  
106 evaluation of predictors of outcome after 30 days from development of severe sepsis or  
107 septic shock due to ESBL-E infection. To perform analysis were created three cohorts:  
108 global, empirical-therapy and targeted-therapy cohorts.

109 **Results:** 367 septic patients were analyzed. Overall mortality was 43.9% at 30 days;  
110 *Escherichia coli* (62.4%) and *Klebsiella pneumoniae* (27.2%) were the most frequent  
111 isolates.  $\beta$ -lactam/ $\beta$ -lactamase inhibitor (BLBLI) combinations was the most  
112 empirically used drug (43.6%), followed by carbapenems (29.4%); empirical therapy  
113 resulted active *in vitro* in 249 (67.8%) patients, but an escalation of antibiotic therapy  
114 was reported in 287 (78.2%) patients. Cox regression analysis showed that age,  
115 Charlson Comorbidity Index, McCabe classification, Pitt bacteremia score, abdominal  
116 source of infection and escalation of antibiotic therapy were independently associated  
117 with 30-day mortality. No differences were reported about survival in patients treated  
118 with BLBLI combinations or carbapenems in empirical or definitive therapy.

119 **Conclusions:** BSI due to ESBL-E in patients who developed severe sepsis or septic  
120 shock was associated with high 30-day mortality; comorbidities, severity scores, source  
121 of infection and need of antibiotic therapy escalation were important determinants of  
122 unfavorable outcome.

**Keywords:** sepsis; septic shock; extended-spectrum  $\beta$ -lactamases; carbapenems;  $\beta$ -lactam/ $\beta$ -lactamase inhibitors

## INTRODUCTION

Bloodstream infections (BSI) caused by extended-spectrum  $\beta$ -lactamases (ESBL)-producing Enterobacteriaceae (E) are associated with high rates of treatment failure and increased mortality, especially when appropriate antimicrobial therapy is delayed [1-2-3-4-5]. On this basis, the choice of an early effective empirical antibiotic therapy in critically ill patients with sepsis and/or septic shock is crucial to reduce the high rates of complications and unfavourable outcome [5].

Previous publications resulting from the INCREMENT project [6-7-8-9-10] highlighted the necessity to identify peculiar clinical and therapeutic features of BSI due to ESBL strains. The role of carbapenems, considered the first choice for the treatment of severe infections caused by ESBL strains [11-12-13-14-15], was redefined also for the high incidence of carbapenem-resistant Enterobacteriaceae strains observed in the last few years [16]. On this basis, attention is now focused on promotion of carbapenem-sparing strategies and evaluation of efficacy of other drugs, like  $\beta$ -lactam/ $\beta$ -lactamase inhibitor combinations (BLBLI), that remain active against a considerable proportion of ESBL-E; however, the role of these drugs is controversial for the treatment of serious infections due to ESBL pathogens [17-18-19].

In this scenario, few data are reported in literature concerning sepsis or septic shock due to ESBL-E infections. Thus, it is important for physicians to recognize peculiar clinical characteristics of sepsis or septic shock due to ESBL-E infections, to promptly identify patients at high risk of unfavorable outcome. Based on this scenario, the aim of this additional study from the INCREMENT project was to assess the predictors of outcome in septic patients with severe sepsis or septic shock caused by ESBL-producing strains.



## MATERIALS AND METHODS

### *Study Design and Patients*

The INCREMENT project is a retrospective international cohort study including patients with clinically significant BSI due to ESBL- or carbapenemase-producing Enterobacteriaceae from January 2004 to December 2013. Characteristics of the INCREMENT study were previously explained [15-16-17-18-19]. This analysis was reported according to the STROBE recommendations [20].

Patients with severe sepsis or septic shock and clinically significant BSI due to ESBL-E were selected from the original database. Data from patients were collected up to 30 days after the diagnosis of BSI from charts; if needed, patients or relatives were contacted by phone.

The INCREMENT project was approved by the Spanish Agency of Medicines (AEMPS; code JRB-ANT-2012-01) and the Hospital Universitario Virgen Macarena Institutional Review Board (code 1921); the need to obtain written informed consent was waived. Approval was also gained at participating centers according to local requirements.

### *Variables, Microbiology and Definitions*

Data collected from all patients included: demographics, acquisition of infection, comorbidities by calculation of Charlson comorbidity index, McCabe classification, Pitt bacteremia score, source of BSI according to clinical and microbiological data, antimicrobial therapy, *in vitro* susceptibility to empirical and targeted antibiotic regimens, length of stay after BSI, and mortality.

Enterobacteriaceae were identified using standard microbiological techniques in each participating center. ESBL-production was screened and confirmed according to CLSI

recommendations [21]; selected isolates from each center had been characterised by polymerase chain reaction (PCR) and DNA sequencing using established methods [17]. Nosocomial acquisition was considered when symptoms of infection started >48 hours after hospital admission or within 48 hours of hospital discharge, while we considered healthcare acquisition if patients had attended haemodialysis or received intravenous chemotherapy in the past 30 days, had been admitted to an acute-care hospital for at least 2 days or had surgery in the past 90 days, or resided in a nursing home or long-term care facility. Other infections were considered community-acquired. For McCabe classification we used three categories: (1) non-fatal (mild and only a few comorbidities), (2) ultimately fatal (risk of death within four years or multiple comorbidities) and (3) rapidly fatal (risk of death during stay, intensive or terminal care patients) underlying diseases. Antimicrobial therapy administered before the susceptibility results were available was considered empirical; therapy administered after microbiological report was considered targeted/definitive. Severe sepsis was defined as sepsis with sepsis-induced organ dysfunction or tissue hypoperfusion (manifesting as hypotension, elevated lactate levels, or decreased urine output); septic shock as severe sepsis plus persistently low blood pressure following the administration of intravenous fluids [22]. Administered BLBLI were amoxicillin-clavulanate (AMC), piperacillin-tazobactam (PTZ), or ampicillin-sulbactam (AMS), quinolones were ciprofloxacin or levofloxacin, aminoglycosides were gentamicin or amikacin, while carbapenems included imipenem, meropenem, and ertapenem. Therapy was considered as monotherapy if no other drugs with intrinsic activity against gram-negative organisms were co-administered, irrespective of isolate susceptibility.

Active empirical therapy was considered the appropriateness of antibiotic therapy for all patients with an etiologic diagnosis according to susceptibility test criteria. We rated an antimicrobial treatment as inadequate if 1 or more of the organisms present were known to have intrinsic resistance or were found to be resistant through susceptibility testing.

Escalation of antibiotic therapy was defined as the switch to or addition of a drug class or classes with a broader spectrum or additional coverage after 48 hours from the initial antibiotic therapy.

### ***Global, Empirical-therapy and Targeted-therapy cohorts***

We constructed 3 non-mutually exclusive cohorts to analyze predictors of mortality, including also therapeutic variables. The global cohort (GC) included all patients analyzed but were explored only pre-treatment variables to identify factors associated with mortality, irrespective of antibiotic therapies. The impact of empirical therapy was investigated in the empirical-therapy cohort (ETC), which included the patients who received a monotherapy that began within the first 24 h after blood cultures were taken and continued for at least 48 h (except for patients who died in  $\leq 48$  h, who were included if they received at least 1 complete day of therapy). The impact of targeted therapy was investigated in the targeted-therapy cohort (TTC), which included the patients who received a monotherapy once the susceptibility profile was available; the targeted drug must have started in  $\leq 5$  days and been administered for at least 50% of the total duration of therapy (except for patients who died while on targeted therapy, who were included if they received at least 1 complete day of therapy).

### ***Endpoints and Statistical Analysis***

The primary endpoint of the study was the evaluation of predictors of outcome at 30 days after development of septic shock due to ESBL infection.

Separate analyses were performed for the three cohorts. To detect significant differences between groups, we used the chi-square test or Fisher exact test for categorical variables, and the 2-tailed  $t$  test or Mann-Whitney test for continuous variables, when appropriate. In univariate and multivariate analysis of survival, the Cox regression model was used to determine the effects of different variables on outcome at 30 days. The cumulative survival was evaluated using Kaplan-Meier product-limit estimators. The final multivariable Cox regression model was selected through forward stepwise regression based on Akaike Information Criterion.

Statistical significance was established at  $\leq 0.05$ . All reported  $P$  values are 2-tailed. The results were obtained using the statistical software R (version 3.3.4; Vienna, Austria).

## RESULTS

The INCREMENT database includes 1005 patients with BSI due to ESBL-E; out of these, 367 (36.5%) patients with severe sepsis or septic shock were analyzed. The number of cases per center in the global cohort ranged from 4 to 50 and wards of hospitalization at the time of ESBL-E isolation, in this study population, are reported in **Figure 1**.

Baseline characteristics and comparison about clinical features of survivors and non-survivors at 30 days are reported in **Table 1**. Overall, mortality at 30 days was observed in 161 (43.9%) patients. Comparison between the two patient groups showed that nosocomial acquisition of infection (64.5% Vs 50%,  $p=0.002$ ), pneumonia (17.4% Vs 8.3%,  $p=0.012$ ), admission in ICU (32.2% Vs 18.9%,  $p=0.004$ ), a higher Charlson Comorbidity index (3 points Vs 2 points,  $p=0.005$ ), a higher Pitt bacteremia score (4 points Vs 3 points,  $<0.001$ ), need of NIV or MV (44% Vs 21.3%,  $p<0.001$ ), and need of vasopressor agents (91.9% Vs 65%,  $p<0.001$ ) were more frequently observed in non-survivors. Conversely, a urinary source of infection (44.7 Vs 24.8,  $p<0.001$ ) was more frequently reported in survivors, compared to patients with poor outcome. Finally, time to appropriate antibiotic therapy (4 days Vs 3 days,  $p=0.01$ ) was longer in non-survivors, compared to survivors.

Etiologies of infection are described in **Table 2**. Overall, *Escherichia coli* (62.4%) and *Klebsiella pneumoniae* (27.2%) were the most frequent isolates; *in vitro* resistance to antibiotics prescribed in empirical therapy, was observed in 4/5 (80%) strains of *Serratia marcescens*, 23/29 (79.3%) of *Enterobacter* spp, 123/229 (53.7%) of *Escherichia coli*, and 41/100 (41%) of *Klebsiella pneumoniae*. Comparison between

survivors and non-survivors showed a more frequent isolation of *Escherichia coli* (67.5% Vs 55.9%,  $p=0.03$ ) in survivors, while *Klebsiella pneumoniae* was more frequently isolated in non-survivors (34.2% Vs 21.8%,  $p=0.012$ ), compared to survivors.

**Table 3** shows a comparison of antibiotics used in empirical and definitive regimens among survivors and non-survivors. BLBLI was the most empirically used class (43.6%), followed by carbapenems (29.4%), and cephalosporins (19.3%), while carbapenems (56.4%), and BLBLI (10.4%) were more frequently used as definitive therapy. A combination therapy was empirically used in 119 (32.4%) patients and in 93 (25.3%) patients as definitive regimen. Empirical antibiotics resulted *in vitro* active in 249 (67.8%) patients, and escalation of antibiotic therapy was used in 287 (78.2%) patients. Comparison between non-survivors and survivors showed that a carbapenem in definitive antibiotic regimen (42.9% Vs 67%,  $p<0.001$ ) and a combination of antibiotics as definitive therapy (16.7% Vs 32%,  $p<0.001$ ) were more frequently used in the latter.

As reported in **Table 4**, univariate and multivariate Cox regression analysis about predictors of outcome at 30 days in the GC showed that age (HR 1.11, CI 95% 1.02-1.22,  $p=0.021$ ), McCabe classification (HR 1.39, CI 95% 1.12-1.72,  $p=0.003$ ), Pitt bacteremia score (HR 1.14, CI 95% 1.09-1.19,  $p<0.001$ ), and abdominal source of infection (HR 1.66, CI 95% 1.08-1.89,  $p=0.02$ ) were independently associated with death.

Based on inclusion criteria (see Methods section), 31 patients were excluded from ETC and TTC to investigate the impact of therapy on 30-day mortality.

The **Table 5** shows that, at univariate and multivariate Cox regression analysis about predictors of outcome at 30 days in 106 patients of the ETC, McCabe classification (HR 1.45, CI 95% 1.14-1.83,  $p=0.002$ ), Pitt bacteremia score (HR 1.14, CI 95% 1.08-1.19,

p<0.001), and escalation of antibiotic therapy (HR 1.9, CI 95% 1.1-3.26, p=0.02) were independently associated with death.

Univariate and multivariate Cox regression analysis about predictors of outcome at 30 days in 230 patients of the TTC is reported in **Table 6**. Charlson Comorbidity Index (HR 1.20, CI 95% 1.11-1.30, p<0.001) and Pitt bacteremia score (HR 1.16, CI 95% 1.09-1.24, p<0.001) were associated with death, while was reported a protective role for quinolones in definitive therapy (HR 0.29, CI 95% 0.11-0.8, p=0.016).

Finally, Kaplan-Meier analysis of 30-day survival of patients in whom an escalation of antibiotic therapy was needed or not is reported in **Figure 2**.

## DISCUSSION

This study highlights the high mortality associated with BSI due to ESBL-E at 30 days from diagnosis in patients who developed severe sepsis or septic shock; furthermore, some clinical characteristics like source of infection (specifically from urinary tract, abdominal or pneumonia), indicators of severity (like comorbidities and admission in ICU), need of respiratory support and/or vasopressor agents were important determinants of outcome in this setting.

However, Cox regression analysis identified age, McCabe classification, Charlson Comorbidity Index, Pitt bacteremia score, abdominal source of infection and escalation of antibiotic therapy (after microbiological report and/or worsening of clinical conditions) as factors independently associated with unfavourable outcome at 30 days. Finally, the use of a quinolone in definitive therapy was associated with 30-day survival.

As previously reported in literature, the severity of clinical conditions (expressed by age, McCabe classification, Charlson Comorbidity Index and Pitt bacteremia score) at time of BSI onset plays a central role in determining the decreased survival in patients with infections due to ESBL-E [23]; furthermore, the progression to sepsis and septic shock is the crucial mechanism associated with high rates of mortality observed in this setting of critically ill patients with BSI due to ESBL strains [24-25-26]. Progression to septic shock may explain the apparently low effectiveness also of an adequate initial



antibiotic therapy, since septic shock is associated with a lethal cascade of events that is unlikely to be interrupted even by an appropriate initial antimicrobial treatment. In addition, most of our patients were severely ill and would probably have been unable to survive their infections independently of the administration of an adequate initial antimicrobial treatment. Finally, the lack of homogeneity in management and treatment of septic patients could partly explain the high rates of mortality.

Of importance, the need of an antibiotic therapy escalation after microbiological report and/or worsening of clinical conditions was independently associated with unfavourable outcome at 30 days, confirming previous observations about an initial inadequate antimicrobial treatment as a major risk factor for mortality [27]. Moreover, severe infections could have an unfavorable outcome despite the administration of adequate antimicrobial treatment, due to the inability to raise plasma drug concentrations above the target MIC. This latter finding may be the cause of underexposure at the infection site, especially in critically ill patients, since septic patients usually require rapid and aggressive fluid resuscitation therapy with a subsequent increase in the extracellular fluid volume that could raise the volume of distribution of drugs [28]. On this basis, differences in rates of mortality could be related to the source of bacteremia. Several studies demonstrated a relationship between source of infection and clinical response [29], with bacteremia secondary to urinary tract infection usually associated with the lowest mortality rate [30]. In our population, as previously reported in literature [29], the abdominal source of infection was an independent determinant of unfavorable outcome considering the abovementioned observations.

An important finding of this analysis was the impact of empirical and definitive antibiotic regimens on 30-day outcome, especially for patients treated with carbapenem or BLBLI. The spread of carbapenemase-producing Enterobacteriaceae promoted

several studies about alternative therapies, like BLBLI. BLBLI has been evaluated for the treatment of these infections [18-31], but no definitive data were reported about the efficacy of BLBLI in the treatment of severe infections, when compared to carbapenems [14-32]. On this basis, the use of carbapenems should be optimized, recommending a role as empirical therapy only for patients who have high risk factors for colonization/infection, based also on geographical patterns of resistance about ESBL strains not susceptible to BLBLI, and on patients who are affected by non-urinary source of bacteremia [33]. In the remaining subjects, in whom clinical conditions or risk factors may not require immediate ESBL antimicrobial coverage, a guided therapy once culture results are obtained may be sufficient [34]. For these reasons, data from international trials will have to assess 30-day mortality associated with the use of BLBLI or carbapenems in the setting of ESBL infections [35]. Of interest, in our analysis quinolones resulted as protective in definitive therapy. The explanation is that, when active *in vitro*, quinolones were a reasonable choice in less critically-ill patients with urinary source of infection. The interpretation of this data is limited by the retrospective design of the study and only randomized trials could address a possible role of quinolones, also in severe infections. However, their use should be carefully evaluated considering the high rates of resistance reported, especially in area with an increased consumption of these antibiotics [36-37].

Of importance, the retrospective nature of the study is a limitation and all the considerations about the impact of empirical or targeted regimens on survival should be done with caution. Nevertheless, our series of septic shock associated with ESBL-E bacteremia is the largest ever reported in literature. Another study limitation is the lack of information of other variables that may influence the outcome, such as source control, circulatory management and PK/PD variables. Finally, no data were recorded

on long-term survival over 30 days; thus, we cannot provide more consistent information on the risk of recurrence of infection and emergence of resistant strains to antibiotic regimen used. However, in the database were applied very strict criteria for the cataloguing of antibiotic regimens and an important strength of this study is the multicenter and international participation to INCREMENT project, with collection of data from different countries.

## CONCLUSIONS

In conclusion, these data point out the high 30-day mortality associated with BSI due to ESBL-E in patients who developed severe sepsis or septic shock, together with the importance of some clinical characteristics, like source of infection and indicators of severity, as determinants of patient's outcome in this setting, requiring an early escalation of antibiotic therapy. Moreover, these data confirm no differences in empirical and definitive antibiotic therapy of BSI, especially for BLBLI and carbapenems, in these critically ill patients. However, randomized clinical trials are needed to confirm these observations.

## ACKNOWLEDGEMENTS

All authors have access to the data and participated in the research and preparation of the manuscript.

## DECLARATIONS

**Funding:** The study was funded by Plan Nacional de I+D+i 2013-2016 and Instituto de Salud Carlos III, Subdirección General de Redes y Centros de Investigación Cooperativa, Ministerio de Economía, Industria y Competitividad, Spanish Network for

Research in Infectious Diseases (REIPI DR12/0015/0010 and RD16/0016/0001) - co-financed by European Development Regional Fund “A way to achieve Europe”, Operative program Intelligent Growth 2014-2020.

**Competing Interests:** None to declare.


**Ethical Approval:** The INCREMENT project was approved by the Spanish Agency of Medicines (AEMPS; code JRB-ANT-2012-01) and the Hospital Universitario Virgen Macarena Institutional Review Board (code 1921).

## APPENDIX

**Other REIPI/ESGBIS/INCREMENT investigators are:** J. Gálvez (Hospital Universitario Virgen Macarena, Seville, Spain); G. Daikos (Laikon General Hospital, Athens, Greece); E. M. Trecarichi and A. R. Losito (Catholic University of the Sacred Heart, Rome, Italy); J. Gómez and E. García-Vázquez (Hospital Universitario Virgen de la Arrixaca, Murcia, Spain); E. Iosifidis and E. Roilides (Hippokration Hospital of Thessaloniki, Thessaloniki, Greece); H. Giamarellou (Hygeia General hospital, Athens, Greece); Y. Doi (University of Pittsburgh, Pittsburgh, USA); F. F. Tuon (Hospital da Universidade Federal do Paraná, Paraná, Brazil); J.A. Martínez, L. Morata and A. Soriano (Hospital Clinic, Barcelona, Spain); F. Navarro and B. Mirelis (Hospital de la Santa Creu i Sant Pau, Barcelona, Spain); R. San Juan and M. Fernández-Ruiz (Hospital 12 de Octubre, Madrid, Spain); N. Larrosa and M. Puig (Hospital Universitario Vall d’Hebrón, Barcelona, Spain); J. Molina and V. González (Hospital Universitario Virgen del Rocío, Seville, Spain); V. Rucci (Hospital Español, Rosario, Argentina); E. Ruiz de Gopegui and C. I. Marinescu (Hospital Universitario Son Espases, Palma de Mallorca, Spain); M. C. Fariñas and M. E. Cano (Hospital Universitario Marqués de Valdecilla-IDIVAL, Santander, Spain); J. R. Paño-Pardo and C. Navarro-San Francisco (Hospital Universitario La Paz-IDIPAZ, Madrid, Spain); S. Gómez-Zorrilla and F. Tubau

472 (Hospital de Bellvitge, Barcelona, Spain); S. Pournaras, A. Tsakris and O. Zarkotou  
473 (University of Athens, Athens, Greece); Ö. K. Azap (Baskent University, Ankara,  
474 Turkey); A. Antoniadou and G. Poulakou (University General Hospital Attikon,  
475 Chiadiri, Greece); D. Virmani (University of Calgary, Calgary, Canada); J. Torre-  
476 Cisneros and J. Guzmán-Puche (Maimonides Biomedical Research Institute of Cordoba  
477 (IMIBIC)/ Reina Sofia University Hospital/University of Cordoba, Córdoba, Spain); Ö.  
478 Helvacı and A. O. Sahin (Hacettepe University, Ankara, Turkey); V. Pintado and R.  
479 Cantón (Hospital Ramón y Cajal, Madrid, Spain); M. Bartoletti and M. Giannella  
480 (Teaching Hospital Policlinico S. Orsola Malpighi, Bologna, Italy); S. Peter (Tübingen  
481 University Hospital, Tübingen, Germany); C. Badia and M. Xercavins (Hospital  
482 Universitario Mútua de Terrassa, Terrassa, Spain); D. Fontanals and E. Jové (Hospital  
483 Parc Taulí, Sabadell, Spain).

484 **Figure 1.** Wards of hospitalization at time of ESBL-E isolation




485

486 **Legend.** ESBL-E: extended-spectrum beta-lactamases-producing Enterobacteriaceae.

487

488 **Figure 2.** Kaplan-Meier curves about 30-day survival of patients in which  
489 was needed or not an escalation of antibiotic therapy\*



490

491 \* $p < 0.001$

492

**Table 1.** Comparison about baseline characteristics and clinical features of survivors and non-survivors at 30 days in the global cohort

Variables	All patients N=367 (%)	Survivors N=206 (%)	Non-Survivors N=161 (%)	<i>p</i>
Age, median (IQR)	69 (57-78)	68 (56-77)	70 (57-80)	0.30
Male sex	216 (58.9)	127 (61.7)	89 (55.3)	0.26
<b>Acquisition of infection</b>				
Community	46 (12.5)	30 (14.5)	16 (9.9)	0.26
Healthcare	106 (28.9)	70 (33.9)	36 (22.3)	<b>0.01</b>
Nosocomial	207 (56.4)	103 (50)	104 (64.5)	<b>0.002</b>
<b>Source of infection</b>				
Primary BSI	22 (6)	10 (4.9)	12 (7.5)	0.41
Urinary	132 (36)	92 (44.7)	40 (24.8)	<b>&lt;0.001</b>
Biliary	36 (9.8)	23 (11.2)	13 (8.1)	0.42
SSTI	7 (1.9)	4 (1.9)	3 (1.9)	1.0
Abdominal	50 (13.6)	22 (10.7)	28 (17.4)	0.09
Pneumonia	45 (12.3)	17 (8.3)	28 (17.4)	<b>0.012</b>
Osteoarticular	2 (0.5)	2 (1)	0	0.59
CNS	1 (0.3)	0	1 (0.6)	0.90
Other	93 (25.3)	60 (29.1)	33 (20.4)	0.085
McCabe classification, nonfatal	145 (39.5)	97 (47)	48 (29.8)	<b>&lt;0.001</b>
Charlson Comorbidity Index, median (IQR)	3 (1-5)	2 (0-4)	3 (2-8)	<b>0.005</b>
Pitt bacteremia score, median (IQR)	3 (2-5)	3 (1-4)	4 (2-7)	<b>&lt;0.001</b>
<b>Comorbidities</b>				
Diabetes	124 (33.8)	60 (29.1)	64 (39.7)	<b>0.027</b>
COPD	77 (21)	37 (17.9)	40 (24.8)	0.09
Myocardial infarct	47 (12.8)	20 (9.7)	27 (16.7)	0.12
Congestive heart failure	70 (19.1)	31 (15)	39 (24.2)	<b>0.027</b>
Peripheral arterial disease	40 (10.9)	16 (7.8)	24 (14.9)	<b>0.046</b>
Dementia	41 (11.2)	22 (10.7)	19 (11.8)	0.83
Immunological disease	16 (4.4)	9 (4.3)	7 (4.3)	1.0
Ulcerative disease	23 (6.3)	12 (5.8)	11 (11.1)	0.81
Liver disease	58 (15.8)	31 (15)	27 (16.7)	0.68
Kidney disease	84 (22.9)	41 (19.9)	43 (26.7)	0.12
Cancer	148 (40.3)	88 (42.7)	60 (37.2)	0.43
Neurological disease	77 (21)	36 (17.4)	41 (25.4)	0.08
AIDS	8 (2.2)	6 (2.9)	2 (1.2)	0.47
ICU admission	91 (24.8)	39 (18.9)	52 (32.2)	<b>0.004</b>
Need of NIV or MV	115 (31.3)	44 (21.3)	71 (44)	<b>&lt;0.001</b>
Need of vasopressor agents	282 (76.8)	134 (65)	148 (91.9)	<b>&lt;0.001</b>



Time to appropriate antibiotic therapy, days, median (IQR)	3 (1-5)	3 (1-4)	4 (2-5)	<b>0.01</b>
--	---------	---------	---------	-------------

495 **Legend.** IQR: inter-quartile range; BSI: bloodstream infection; SSTI: skin and soft tissue infection; CNS:  
 496 central nervous system; COPD: chronic obstructive pulmonary disease; AIDS: acquired immune  
 497 deficiency syndrome; ICU: intensive care unit; NIV: non-invasive mechanical ventilation; MV:  
 498 mechanical ventilation.

499

500

501 **Table 2.** Etiology of infection among survivors and non-survivors at 30  
 502 days and rate of resistance to empirical therapy

Pathogens	Survivors N=206 (%)	Non-Survivors N=161 (%)	<i>p</i>	Resistance to empirical therapy N° resistant isolates/ total isolates (%)
<i>Escherichia coli</i>	139 (67.5)	90 (55.9)	<b>0.03</b>	123/229 (53.7)
<i>Klebsiella pneumoniae</i>	45 (21.8)	55 (34.2)	<b>0.012</b>	41/100 (41)
<i>Enterobacter</i> spp	17 (8.3)	12 (7.5)	0.93	23/29 (79.3)
<i>Serratia marcescens</i>	3 (1.5)	2 (1.2)	1.0	4/5 (80)
<i>Proteus</i> spp	0	2 (1.2)	0.37	0
<i>Citrobacter</i> spp	1 (0.5)	0	1.0	0
<i>Morganella morganii</i>	1 (0.5)	0	1.0	0

503

504 **Table 3.** Comparison between antibiotics administered in empirical and  
 505 definitive therapy to survivors and non-survivors at 30 days in the global  
 506 cohort

Antibiotics	Empirical therapy			Definitive therapy		
	Survivors N=206 (%)	Non-Survivors N=161 (%)	<i>p</i>	Survivors N=206 (%)	Non-Survivors N=161 (%)	<i>p</i>
Cephalosporin	43 (20.9)	28 (17.4)	0.48	3 (1.5)	1 (0.6)	0.80
BLBLI	94 (45.6)	66 (41)	0.43	27 (13.1)	11 (6.8)	0.07

Aminoglycosides	22 (10.7)	21 (13)	0.59	16 (7.8)	8 (5)	0.39
Quinolones	30 (14.6)	25 (15.5)	0.91	14 (6.8)	6 (3.7)	0.29
TMP/SMX	1 (0.5)	0	1.0	8 (3.9)	4 (2.5)	0.65
Fosfomycin	2 (1)	0	0.59	0	0	-
Tetracycline	2 (1)	2 (1.2)	1.0	0	0	-
Monobactams	1 (0.5)	1 (0.6)	1.0	0	0	-
Colistin	1 (0.5)	1 (0.6)	1.0	3 (1.5)	1 (0.6)	0.79
Carbapenem	61 (29.6)	47 (29.2)	0.44	138 (67)	69 (42.9)	<0.001
Tigecycline	0	0	-	6 (2.9)	2 (1.2)	0.47
Chloramphenicol	0	0	-	1 (0.5)	0	1.0
Combination therapy	150 (72.8)	111 (68.9)	0.54	88 (42.7)	49 (30.4)	0.001
Median length of therapy (IQR)	3 (1-4)	2 (1-4)	0.1	11 (5-14)	7 (3-10)	<0.001
Active empirical therapy	143 (69.4)	106 (65.8)	0.54			
Escalation of antibiotic therapy	154 (74.7)	133 (82.6)	0.09			

507 **Legend.** BLBLI:  $\beta$ -lactam/ $\beta$ -lactamase inhibitor; TMP/SMX: Trimethoprim/sulfamethoxazole; IQR:  
508 inter-quartile range.

509 **Note.** BLBLI: amoxicillin/clavulanate or ampicillin/sulbactam or piperacillin-tazobactam

510 Quinolones: ciprofloxacin or levofloxacin

511 Aminoglycosides: gentamicin or amikacin

512 Carbapenem: meropenem or imipenem or ertapenem

513

514 **Table 4.** Univariate and Multivariate Cox regression analysis about  
515 predictors of outcome at 30 days in 367 patients of the global cohort

516

VARIABLES	UNIVARIATE			MULTIVARIATE		
	HR	95%CI	<i>p</i>	HR	95%CI	<i>p</i>
Age	1.1	1.01-1.21	0.031	1.11	1.02-1.22	0.021
Male sex	0.84	0.62-1.15	0.284			
<b>Acquisition of infection</b>						
Healthcare	0.95	0.53-1.71	0.863			
Nosocomial	1.38	0.81-2.34	0.233			

<b>Source of infection</b>						
Primary BSI	1.08	0.6-1.94	0.807			
Urinary	0.59	0.41-0.84	0.003			
Biliary	0.81	0.46-1.44	0.478			
SSTI	0.82	0.26-2.57	0.735			
Abdominal	1.46	0.97-2.19	0.07	1.66	1.08-1.89	0.02
Pneumonia	0.45	1.04-2.36	0.3			
ICU admission	1.4	1.01-1.95	0.047			
Charlson Comorbidity Index	1.05	0.99-1.11	0.083			
McCabe classification	1.54	1.25-1.89	<0.001	1.39	1.12-1.72	0.003
Pitt bacteremia score	1.14	1.09-1.19	<0.001	1.14	1.09-1.19	<0.001
<b>Comorbidities</b>						
Diabetes	1.42	1.03-1.95	0.033			
COPD	1.28	0.89-1.84	0.175			
Myocardial infarct	1.29	0.85-1.96	0.238			
Congestive heart failure	1.28	0.87-1.84	0.19			
Peripheral arterial disease	1.41	0.91-2.18	0.12			
Dementia	1.08	0.67-1.75	0.752			
Immunological disease	0.82	0.39-1.76	0.617			
Ulcerative disease	1.08	0.58-1.99	0.816			
Liver disease	1.11	0.73-1.68	0.629			
Kidney disease	1.31	0.92-1.87	0.132			
Cancer	0.85	0.62-1.18	0.334			

AIDS	0.43	0.11-1.72	0.232			
Need of NIV or MV	1.93	1.4-2.64	<0.001			
Need of vasopressor agents	1.37	1.01-2.08	0.044			

517 **Legend.** HR: hazard ratio; CI: confidence interval; BSI: bloodstream infection; SSTI: skin and soft tissue  
518 infection; ICU: intensive care unit; COPD: chronic obstructive pulmonary disease; AIDS: acquired  
519 immune deficiency syndrome; NIV: non-invasive mechanical ventilation; MV: mechanical ventilation.

520

521 **Table 5.** Univariate and Multivariate Cox regression analysis about  
522 predictors of outcome at 30 days in 106 patients of the empirical-therapy  
523 cohort

524

VARIABLES	UNIVARIATE			MULTIVARIATE		
	HR	CI 95%	<i>p</i>	HR	CI 95%	<i>p</i>
Age	1.09	0.99-1.19	0.075			
Male sex	0.82	0.59-1.13	0.226			
<b>Acquisition of infection</b>						
Healthcare	0.95	0.52-1.74	0.861			
Nosocomial	1.36	0.79-2.35	0.270			
<b>Source of infection</b>						
Primary BSI	1.19	0.66-2.15	0.562			
Urinary	0.56	0.39-0.82	0.003			
Biliary	0.87	0.49-1.54	0.639			
SSTI	0.85	0.27-2.68	0.786			
Abdominal	1.42	0.93-2.17	0.104			
Pneumonia	1.79	1.18-2.72	0.006			

ICU admission	1.58	1.12-2.23	0.009			
Charlson Comorbidity Index	1.06	0.99-1.12	0.069			
McCabe classification	1.55	1.25-1.93	<0.001	1.45	1.14-1.83	0.002
Pitt bacteremia score	1.16	1.11-1.22	<0.001	1.14	1.08-1.19	<0.001
<b>Comorbidities</b>						
Diabetes	1.37	0.98-1.92	0.062			
COPD	1.45	1.01-2.1	0.047			
Myocardial infarct	1.43	0.92-2.21	0.111			
Congestive heart failure	1.43	0.98-2.09	0.06			
Peripheral arterial disease	1.25	0.77-2.03	0.357			
Dementia	0.98	0.58-1.65	0.934			
Immunological disease	0.79	0.35-1.80	0.58			
Ulcerative disease	1.09	0.57-2.07	0.797			
Liver disease	1.12	0.73-1.73	0.593			
Kidney disease	1.43	0.99-2.05	0.056			
Cancer	0.91	0.65-1.27	0.578			
AIDS	0.44	0.11-1.78	0.249			
Need of NIV or MV	1.96	1.41-2.72	<0.001			
Need of vasopressor agents	1.69	1.14-2.49	0.008			
Cephalosporin	0.87	0.57-1.31	0.494			
BLBLI	0.95	0.69-1.32	0.763			
Aminoglycosides	1.06	0.67-1.68	0.808			
Quinolones	1.12	0.73-1.73	0.595			

Tetracycline	1.34	0.33-5.44	0.677			
Carbapenem	1.02	0.72-1.44	0.914			
Combination therapy	0.78	0.65-1.11	0.12			
Escalation of antibiotic therapy	1.92	1.12-3.28	0.017	1.9	1.1-3.26	0.02
Time to appropriate antibiotic therapy	1.2	0.87-2.34	0.25			

525 **Legend.** HR: hazard ratio; CI: confidence interval; BSI: bloodstream infection; SSTI: skin and soft tissue  
526 infection; ICU: intensive care unit; COPD: chronic obstructive pulmonary disease; AIDS: acquired  
527 immune deficiency syndrome; NIV: non-invasive mechanical ventilation; MV: mechanical ventilation.

528

529 **Table 6.** Univariate and Multivariate Cox regression analysis about  
530 predictors of outcome at 30 days in 230 patients of the targeted-therapy  
531 cohort

532

VARIABLES	UNIVARIATE			MULTIVARIATE		
	HR	CI 95%	<i>p</i>	HR	CI 95%	<i>p</i>
Age	1.1	1.01-1.21	0.031			
Male sex	1.19	0.62-1.15	0.284			
<b>Acquisition of infection</b>						
Healthcare	0.95	0.53-1.72	0.863			
Nosocomial	1.38	0.81-2.34	0.233			
<b>Source of infection</b>						
Primary BSI	1.73	0.79-3.76	0.169			
Urinary	0.68	0.41-1.13	0.137			
Biliary	1.53	0.79-2.99	0.208			

SSTI	1.71	0.54-5.41	0.365			
Abdominal	0.92	0.44-1.92	0.829			
Pneumonia	1.52	0.87-2.68	0.144			
ICU admission	1.4	1.01-1.95	0.047			
Charlson Comorbidity Index	1.05	0.99-1.11	0.083	1.20	1.11-1.30	<0.001
McCabe classification	1.54	1.25-1.89	<0.001			
Pitt bacteremia score	1.14	1.09-1.19	<0.001	1.16	1.09-1.24	<0.001
<b>Comorbidities</b>						
Diabetes	1.42	1.03-1.95	0.033			
COPD	1.29	0.89-1.84	0.175			
Myocardial infarct	1.29	0.85-1.96	0.238			
Congestive heart failure	1.28	0.89-1.84	0.19			
Peripheral arterial disease	1.41	0.91-2.18	0.12			
Dementia	1.08	0.67-1.75	0.752			
Immunological disease	0.66	0.21-2.09	0.475			
Ulcerative disease	1.08	0.58-1.99	0.816			
Liver disease	1.11	0.73-1.68	0.629			
Kidney disease	1.31	0.92-1.87	0.132			
Cancer	0.85	0.62-1.18	0.334			
AIDS	0.43	0.11-1.72	0.232			
Need of NIV or MV	1.88	1.18-3.0	0.008			
Need of vasopressor agents	1.32	0.79-2.18	0.287			
Cephalosporin	1.46	0.1-4.91	0.708			

BLBLI	1.64	0.33-1.13	0.114			
Aminoglycosides	0.59	0.29-1.20	0.142			
Quinolones	0.54	0.24-1.22	0.141	0.29	0.11-0.8	0.016
TMP/SMX	1.01	0.32-3.19	0.992			
Colistin	0.5	0.07-3.61	0.493			
Carbapenem	0.96	0.54-1.71	0.883			
Tigecycline	0.3	0.04-2.2	0.239			
Combination therapy	0.59	0.39-1.22	0.24			

**Legend.** HR: hazard ratio; CI: confidence interval; BSI: bloodstream infection; SSTI: skin and soft tissue infection; ICU: intensive care unit; COPD: chronic obstructive pulmonary disease; AIDS: acquired immune deficiency syndrome; NIV: non-invasive mechanical ventilation; MV: mechanical ventilation.



554 **REFERENCES**

- 
- <sup>1</sup> Rodríguez-Baño J, Navarro MD, Romero L, Muniain MA, de Cueto M, Ríos MJ, et al. Bacteremia due to extended-spectrum beta-lactamase-producing *Escherichia coli* in the CTX-M era: a new clinical challenge. Clin Infect Dis 2006;43:1407-14.
- <sup>2</sup> Kang CI, Kim SH, Park WB, Lee KD, Kim HB, Kim EC, et al. Bloodstream infections caused by antibiotic-resistant gram negative bacilli: risk factors for mortality and impact of inappropriate initial antimicrobial therapy on outcome. Antimicrob Agents Chemother 2005;49:760-6.
- <sup>3</sup> Peralta G, Sanchez MB, Garrido JC, De Benito I, Cano ME, Martínez-Martínez L, et al. Impact of antibiotic resistance and of adequate empirical antibiotic treatment in the prognosis of patients with *Escherichia coli* bacteraemia. J Antimicrob Chemother 2007;60:855-63.
- <sup>4</sup> Ortega M, Marco F, Soriano A, Almela M, Martínez JA, Muñoz A, et al. Analysis of 4758 *Escherichia coli* bacteraemia episodes: predictive factors for isolation of an antibiotic-resistant strain and their impact on the outcome. J Antimicrob Chemother 2009;63:568-74.
- <sup>5</sup> Tumbarello M, Sanguinetti M, Montuori E, Trecarichi EM, Posteraro B, Fiori B, et al. Predictors of mortality in patients with bloodstream infections caused by extended-spectrum beta-lactamase-producing Enterobacteriaceae: importance of inadequate initial antimicrobial treatment. Antimicrob Agents Chemother 2007;51:1987-94.
- <sup>6</sup> Gutiérrez-Gutiérrez B, Salamanca E, de Cueto M, Hsueh PR, Viale P, Paño-Pardo JR, et al; REIPI/ESGBIS/INCREMENT Investigators. Effect of appropriate combination therapy on mortality of patients with bloodstream infections due to carbapenemase-producing Enterobacteriaceae (INCREMENT): a retrospective cohort study. Lancet Infect Dis. 2017;17:726-734.
- <sup>7</sup> Palacios-Baena ZR, Gutiérrez-Gutiérrez B, De Cueto M, Viale P, Venditti M, Hernández-Torres A, et al; REIPI/ESGBIS/INCREMENT Group. Development and validation of the INCREMENT-ESBL

predictive score for mortality in patients with bloodstream infections due to extended-spectrum- $\beta$ -lactamase-producing Enterobacteriaceae. *J Antimicrob Chemother*. 2017;72:906-913.

<sup>8</sup> Gutiérrez-Gutiérrez B, Salamanca E, de Cueto M, Hsueh PR, Viale P, Paño-Pardo JR, et al; Investigators from the REIPI/ESGBIS/INCREMENT Group. A predictive model of mortality in patients with bloodstream infections due to carbapenemase-producing Enterobacteriaceae. *Mayo Clin Proc*. 2016;91:1362-1371.

<sup>9</sup> Gutiérrez-Gutiérrez B, Pérez-Galera S, Salamanca E, de Cueto M, Calbo E, Almirante B, et al. A multinational, preregistered cohort study of  $\beta$ -Lactam/ $\beta$ -Lactamase inhibitor combinations for treatment of bloodstream infections due to extended-spectrum- $\beta$ -Lactamase-producing Enterobacteriaceae. *Antimicrob Agents Chemother*. 2016;60:4159-69.

<sup>10</sup> Gutiérrez-Gutiérrez B, Bonomo RA, Carmeli Y, Paterson DL, Almirante B, Martínez-Martínez L, et al; REIPI/ESGBIS/INCREMENT Group. Ertapenem for the treatment of bloodstream infections due to ESBL-producing Enterobacteriaceae: a multinational pre-registered cohort study. *J Antimicrob Chemother*. 2016;71:1672-80.

<sup>11</sup> Paterson DL, Bonomo R. Extended-spectrum  $\beta$ -lactamase: a clinical update. *Clin Microbiol Rev* 2005;18:657–86.

<sup>12</sup> Pitout JDD, Laupland KB. Extended-spectrum  $\beta$ -lactamase-producing Enterobacteriaceae: an emerging public-health concern. *Lancet Infect Dis* 2008;8:159–66.

<sup>13</sup> Rodríguez-Baño J, Pascual A. Clinical significance of extended-spectrum  $\beta$ -lactamases. *Expert Rev Anti Infect Ther* 2008;6:671–83.

<sup>14</sup> Schwaber MJ, Carmeli Y. Carbapenem-resistant Enterobacteriaceae. A potential threat. *JAMA* 2008;300:2911–3.

<sup>15</sup> Hawser S, Hoban D, Bouchillon S, Badal R, Carmeli Y, Hawkey P. Antimicrobial susceptibility of intra-abdominal gram-negative bacilli from Europe: SMART Europe 2008. *Eur J Clin Microbiol Infect Dis* 2011;30:173–9.

- <sup>16</sup> Orsi GB, Falcone M, Venditti M. Surveillance and management of multidrug-resistant microorganisms. *Expert Rev Anti Infect Ther* 2011; 9: 653-79.
- <sup>17</sup> Rodríguez-Baño J, Navarro MD, Retamar P, Picón E, Pascual Á; Extended-Spectrum Beta-Lactamases–Red Española de Investigación en Patología Infecciosa/Grupo de Estudio de Infección Hospitalaria Group.  $\beta$ -Lactam/ $\beta$ -lactam inhibitor combinations for the treatment of bacteremia due to extended-spectrum  $\beta$ -lactamase–producing *Escherichia coli*: a post hoc analysis of prospective cohorts. *Clin Infect Dis* 2012; 54:167–74.
- <sup>18</sup> Vardakas KL, Tansarli GS, Rafailidis PI, Falagas ME. Carbapenems versus alternative antibiotics for the treatment of bacteremia due to Enterobacteriaceae producing extended-spectrum  $\beta$ -lactamases: a systematic review and meta-analysis. *J Antimicrob Chemother.* 2012; 67:2793-803.
- <sup>19</sup> Tamma PD, Han JH, Rock C, Harris AD, Lautenbach E, Hsu AJ, et al. Carbapenem therapy is associated with improved survival compared with piperacillin-tazobactam for patients with extended-spectrum  $\beta$ -lactamase bacteremia. *Clin Infect Dis* 2015;60:1319-25.
- <sup>20</sup> von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP; STROBE Initiative. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet* 2007; 370:1453–7.
- <sup>21</sup> Clinical and Laboratory Standard Institute. Performance standards for antimicrobial susceptibility testing. CLSI M100-S22. CLSI, 2012.
- <sup>22</sup> Dellinger RP, Levy MM, Rhodes A, Annane D, Gerlach H, Opal SM, et al; Surviving Sepsis Campaign Guidelines Committee including the Pediatric Subgroup. Surviving Sepsis Campaign: international guidelines for management of severe sepsis and septic shock, 2012. *Intensive Care Med.* 2013;39:165-228.
- <sup>23</sup> Cosgrove SE. The relationship between antimicrobial resistance and patient outcomes: mortality, length of hospital stay, and health care costs. *Clin Infect Dis* 2006;42:S82-9.

- <sup>24</sup> Metan G, Zarakolu P, Cakir B, Hascelik G, Uzun O. Clinical outcomes and therapeutic options of bloodstream infections caused by extended spectrum beta-lactamase-producing *Escherichia coli*. Int J Antimicrob Agents 2005; 26: 254-7.
- <sup>25</sup> Lautenbach E, Patel JB, Bilker WB, Edelstein PH, Fishman NO. Extended-spectrum betalactamase-producing *Escherichia coli* and *Klebsiella pneumoniae*: risk factors for infection and impact of resistance on outcomes. Clin Infect Dis 2001;32:1162-71.
- <sup>26</sup> Kang CI, Kim SH, Kim DM, Park WB, Lee KD, Kim HB, et al. Risk factors for and clinical outcomes of bloodstream infections caused by extended-spectrum betalactamase-producing *Klebsiella pneumoniae*. Infect Control Hosp Epidemiol 2004;25:860-7.
- <sup>27</sup> Garnacho-Montero J, Gutiérrez-Pizarra A, Escoreca-Ortega A, Corcia-Palomo Y, Fernández-Delgado E, Herrera-Melero I, et al. De-escalation of empirical therapy is associated with lower mortality in patients with severe sepsis and septic shock. Intensive Care Med. 2014;40:32-40
- <sup>28</sup> Pea F, Viale P. Bench-to-bedside review: Appropriate antibiotic therapy in severe sepsis and septic shock--does the dose matter? Crit Care. 2009;13:214.
- <sup>29</sup> Falcone M, Russo A, Iacovelli A, Restuccia G, Ceccarelli G, Giordano A, et al. Predictors of outcome in ICU patients with septic shock caused by *Klebsiella pneumoniae* carbapenemase-producing *K. pneumoniae*. Clin Microbiol Infect. 2016;22:444-50.
- <sup>30</sup> Paterson DL, Ko WC, Von Gottberg A, Mohapatra S, Casellas JM, Goossens H, et al. Antibiotic therapy for *Klebsiella pneumoniae* bacteremia: implications of production of extended-spectrum beta-lactamases. Clin Infect Dis 2004;39:31-7.
- <sup>31</sup> Shiber S, Yahav D, Avni T, Leibovici L, Paul M.  $\beta$ -Lactam/ $\beta$ -lactamase inhibitors versus carbapenems for the treatment of sepsis: systematic review and meta-analysis of randomized controlled trials. J Antimicrob Chemother. 2015; 70:41-7.
- <sup>32</sup> López-Cerero L, Picón E, Morillo C, Hernández JR, Docobo F, Pachón J, et al. Comparative assessment of inoculum effects on the antimicrobial activity of amoxicillin/clavulanate and

piperacillin/tazobactam with ESBL-producing and non-producing *Escherichia coli* isolates. Clin Microbiol Infect 2010; 16:132–6.

<sup>33</sup> Harris PNA, Pezzani MD, Gutiérrez-Gutiérrez B, Viale P, Hsueh PR, Ruiz-Garbajosa P, et al; ESGBI/REIPI/INCREMENT group. Geographical variation in therapy for bloodstream infections due to multidrug-resistant enterobacteriaceae: a post hoc analysis of the INCREMENT study. Int J Antimicrob Agents. 2017;50:664-672

<sup>34</sup> Tumbarello M, Trecarichi EM, Bassetti M, Bassetti M, De Rosa FG, Spanu T, Di Meo E, et al. Identifying patients harboring extended-spectrum-beta-lactamase-producing Enterobacteriaceae on hospital admission: derivation and validation of a scoring system. Antimicrob Agents Chemother 2011;55:3485-90.

<sup>35</sup> Harris PN, Peleg AY, Iredell J, Ingram PR, Miyakis S, Stewardson AJ, et al. Meropenem versus piperacillin-tazobactam for definitive treatment of bloodstream infections due to ceftriaxone non-susceptible *Escherichia coli* and *Klebsiella* spp (the MERINO trial): study protocol for a randomised controlled trial. Trials. 2015 Jan 27;16:24.

<sup>36</sup> Sarma JB, Marshall B, Cleeve V, Tate D, Oswald T, Woolfrey S. Effects of fluoroquinolone restriction (from 2007 to 2012) on resistance in Enterobacteriaceae: interrupted time-series analysis. J Hosp Infect. 2015;91:68-73.

<sup>37</sup> Cha MK, Kang CI, Kim SH, Cho SY, Ha YE, Wi YM, et al; Korean Network for Study on Infectious Diseases (KONSID). In vitro activities of 21 antimicrobial agents alone and in combination with aminoglycosides or fluoroquinolones against extended-spectrum- $\beta$ -lactamase-producing *Escherichia coli* isolates causing bacteremia. Antimicrob Agents Chemother. 2015;59:5834-7.